



HALF-FRASER BROTH BASE, 12.9 g (G164-12.9) SampleReady® GAMMA IRRADIATED SOLUBLE POUCH

USE: Half-Fraser Broth Base with Supplement is used for the selective enrichment of *Listeria monocytogenes*.

DESCRIPTION: Half Fraser Broth is a modification of Fraser Broth which contains half of the concentration of nalidixic acid and acriflavine hydrochloride to aid in the recovery of stressed cells. Half Fraser Broth is used as the primary enrichment broth in the ISO methodology⁵ for the detection of *Listeria*.

Fraser Broth Base and Fraser Broth Supplement are based on the Fraser Broth formulation of Fraser and Sperber.¹ The medium is used in the rapid detection of *Listeria* from food and environmental samples. Demi-Fraser Broth Base is a modification of Fraser Broth Base in which the nalidixic acid and acriflavine concentrations have been reduced to 10 mg/L and 12.5 mg/L respectively.² Peptone, beef extract, and yeast extract provide carbon and nitrogen sources and the cofactors required for good growth of *Listeria*. Sodium phosphate and potassium phosphate buffer the medium. Selectivity is provided by lithium chloride, nalidixic acid, and acriflavine. The high sodium chloride concentration of the medium inhibits growth of enterococci. All *Listeria* species hydrolyze esculin, as evidenced by a blackening of the medium. This blackening results from the formation of 6,7 dihydroxycoumarin, which reacts with the ferric ions.¹ Ferric ions are added to the final medium as ferric ammonium citrate in Fraser Broth Supplement. Some molecular detection methodologies do not require supplementation with Ferric Ammonium Citrate.

FORMULA:

Proteose Peptone.....	5.0 g/L
Beef Extract.....	5.0 g/L
Meat Peptone.....	5.0 g/L
Casein Peptone.....	5.0 g/L
Sodium Chloride.....	20.0 g/L
Disodium Phosphate.....	12.0 g/L
Monopotassium Phosphate.....	1.35 g/L
Esculin.....	1.0 g/L
Nalidixic Acid.....	10 mg/L
Acriflavine HCl.....	12.5 mg/L
Lithium Chloride.....	3.0 g/L
Total.....	57.37 g/L

Note: Medium may be adjusted and/or supplemented as required to meet performance criteria.

Fraser Broth Supplement:

Per 10mL Vial	
Ferric Ammonium Citrate.....	0.5 g/L

Final pH: 7.2 ± 0.2 at 25°C

PHYSICAL APPEARANCE:

Dehydrated Appearance – The powder is tan, homogenous, and free flowing encapsulated in a clear soluble film pouch.

Prepared Appearance – Medium amber, clear to slightly opalescent with a fine precipitate.

PROCEDURE:

1. Carefully open the Mylar bag and aseptically transfer one soluble pouch to a container with 225 ml sterile water and mix.
2. Dissolve completely with repeated stirring or agitation.
3. Once dissolved, the medium is ready for testing applications. Aseptically add Fraser Broth Supplement.
4. Consult reference methods for complete procedures.

EXPECTED RESULTS: Cultural response after 18-48 hours at 35°C.

Microorganism	CFU	Growth	Esculin Reaction
<i>E. faecalis</i> ATCC™ 29212	10 ³ -2x10 ³	-	-
<i>E. coli</i> ATCC™ 25922	10 ³ -2x10 ³	-	-
<i>L. monocytogenes</i> ATCC™ 19114	10 ² -10 ³	+	blackening
<i>S. aureus</i> ATCC™ 25923	10 ³ -2x10 ³	-	-

STORAGE: Store the sealed Mylar bag at 2-30°C in a dry environment for up to the expiration date.

LIMITATIONS: Once opened, use all pouches within the Mylar bag as soon as possible. Use prepared media within 3 hours for best results. The pouches should be discarded if there has been a change from the original color, or the encapsulated powder is not free flowing.

For laboratory use only.

SIZES AVAILABLE: 5.2 g (90 ml), 12.9 g (225 ml) 29 g (500 ml)

PACKAGING: One box contains a total of 300 pouches consisting of 60 hermetically sealed Mylar bags containing 5 soluble pouches containing 12.9 g of dehydrated culture media each. Additional configurations are available upon request.

REFERENCES:

1. Fraser J.A. and Sperber W.H. 1988. J. Food Protect. 51, No.10, 762-765.
2. McClain D. and Lee W.H. 1988. J. Assoc. Off. Ana. Chem. 71, No.3, 660-664.
3. Cowart R.E. and Foster B.G. 1985. J. Infect. Dis. 151, 721-730.
4. Partis L., Newton K., Marby J. and Wells R.J. 1994. Appl. Env. Microbiol. 60, 1693-1694.
5. Microbiology of Food and Animal Feeding Stuffs – Horizontal method for the detection and enumeration of *Listeria monocytogenes* Part 1: Detection Method. BS EN ISO 11290:1 1997.